

**Rehospitalization Intervention
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies



PEL/VIP Corporate Total
Cardio-Pulmonary Participant Report
July 2017

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- 17.1% for Pneumonia
- 21.9% for CHF
- 20.0% for COPD


*medicare.gov/hospitalcompare

Results Summary: 1684 Total Participants

Participant Breakdown: Pneumonia - 659 High Risk, 506 Low Risk; CHF - 666 High Risk, 512 Low Risk; COPD - 365 High Risk, 130 Low Risk

Re-Hospitalization Results

0.8% Pneumonia
1.1% CHF
0.2% COPD



30-Day Nursing Home CMS Quality Measures

- 25% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 9% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*500 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program