



**Rehospitalization Intervention
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- 17.1% for Pneumonia
- 21.9% for CHF
- 20.0% for COPD

PEL/VIP Whole Company Outcomes
Cardio-Pulmonary Participant Report
September 2017


*medicare.gov/hospitalcompare

Results Summary: 1860 Total Participants

Participant Breakdown: Pneumonia - 644 High Risk, 550 Low Risk; CHF - 851 High Risk, 504 Low Risk; COPD - 410 High Risk, 130 Low Risk

Re-Hospitalization Results

0.6% Pneumonia
0.4% CHF
0.6% COPD



30-Day Nursing Home CMS Quality Measures

- 23% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 8% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*532 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program