

**Rehospitalization Intervention  
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies



**PEL/VIP Corporate**  
Cardio-Pulmonary Participant Report  
December 2017

**The U.S. Average for 30-Day  
Re-Hospitalization Rates\*:**

- 17.1% for Pneumonia
- 21.9% for CHF
- 20.0% for COPD


\*medicare.gov/hospitalcompare

**Results Summary: 1852 Total Participants**

Participant Breakdown: Pneumonia - 772 High Risk, 518 Low Risk; CHF - 992 High Risk, 461 Low Risk; COPD - 441 High Risk, 96 Low Risk

Re-Hospitalization Results

0.9% Pneumonia  
0.9% CHF  
0.7% COPD



30-Day Nursing Home CMS Quality Measures

- 22% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 10% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*604 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program