

**Rehospitalization Intervention
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies



PEL/VIP Company Wide
Cardio-Pulmonary Participant Report
January 2018

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- 16.9% for Pneumonia
- 21.6% for CHF
- 19.8% for COPD


*medicare.gov/hospitalcompare

Results Summary: 1963 Total Participants

Participant Breakdown: Pneumonia - 921 High Risk, 461 Low Risk; CHF - 1054 High Risk, 451 Low Risk; COPD - 475 High Risk, 113 Low Risk

Re-Hospitalization Results

1.5% Pneumonia
0.6% CHF
0.7% COPD



30-Day Nursing Home CMS Quality Measures

- 22% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 11% of short stay participants were re-hospitalized for any reason
- 3% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*587 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program