



**Rehospitalization Intervention  
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies

**The U.S. Average for 30-Day  
Re-Hospitalization Rates\*:**

- 16.9% for Pneumonia
- 21.6% for CHF
- 19.8% for COPD

**PEL/VIP Whole Company**  
Cardio-Pulmonary Participant Report  
April 2018


\*medicare.gov/hospitalcompare

**Results Summary: 1723 Total Participants**

Participant Breakdown: Pneumonia - 794 High Risk, 429 Low Risk; CHF - 955 High Risk, 354 Low Risk; COPD - 448 High Risk, 82 Low Risk

Re-Hospitalization Results

0.6% Pneumonia  
0.7% CHF  
0.8% COPD



30-Day Nursing Home CMS Quality Measures

- 23% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 10% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*478 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program