



**Rehospitalization Intervention
Program Objectives:**

- . Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- . Initiate Respiratory Therapy into the Plan of Care
- . Utilize Prevention Strategies

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- . 16.9% for Pneumonia
- . 21.6% for CHF
- . 19.8% for COPD

PEL/VIP Whole Company
Cardio-Pulmonary Participant Report
June 2018


*medicare.gov/hospitalcompare

Results Summary: 1521 Total Participants

Participant Breakdown: Pneumonia - 679 High Risk, 378 Low Risk; CHF - 871 High Risk, 330 Low Risk; COPD - 389 High Risk, 75 Low Risk

Re-Hospitalization Results

0.8% Pneumonia
0.5% CHF
0.6% COPD



30-Day Nursing Home CMS Quality Measures

- 25% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 10% of short stay participants were re-hospitalized for any reason
- 3% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*324 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program