

**Rehospitalization Intervention  
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies



**PEL/VIP Whole Company**  
Cardio-Pulmonary Participant Report  
October 2018

**The U.S. Average for 30-Day  
Re-Hospitalization Rates\*:**

- 16.7% for Pneumonia
- 21.7% for CHF
- 19.6% for COPD


\*medicare.gov/hospitalcompare

**Results Summary: 1082 Total Participants**

Participant Breakdown: Pneumonia - 561 High Risk, 104 Low Risk; CHF - 712 High Risk, 64 Low Risk; COPD - 398 High Risk, 16 Low Risk

Re-Hospitalization Results

0.6% Pneumonia  
0.5% CHF  
0.2% COPD



30-Day Nursing Home CMS Quality Measures

- 24% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 12% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*420 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program