

**Rehospitalization Intervention
Program Objectives:**

- . Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- . Initiate Respiratory Therapy into the Plan of Care
- . Utilize Prevention Strategies



PEL/VIP Whole Company
Cardio-Pulmonary Participant Report
November 2018

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- . 16.7% for Pneumonia
- . 21.7% for CHF
- . 19.6% for COPD


*medicare.gov/hospitalcompare

Results Summary: 1120 Total Participants

Participant Breakdown: Pneumonia - 559 High Risk, 87 Low Risk; CHF - 725 High Risk, 57 Low Risk; COPD - 452 High Risk, 23 Low Risk

Re-Hospitalization Results

2.0% Pneumonia
1.0% CHF
0.8% COPD



30-Day Nursing Home CMS Quality Measures

- 22% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 11% of short stay participants were re-hospitalized for any reason
- 4% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*446 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program