

**Rehospitalization Intervention
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies



PEL/VIP Whole Company
Cardio-Pulmonary Participant Report
January 2019

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- 16.7% for Pneumonia
- 21.7% for CHF
- 19.6% for COPD


*medicare.gov/hospitalcompare

Results Summary: 1254 Total Participants

Participant Breakdown: Pneumonia - 599 High Risk, 67 Low Risk; CHF - 817 High Risk, 50 Low Risk; COPD - 479 High Risk, 16 Low Risk

Re-Hospitalization Results

1.4% Pneumonia
1.0% CHF
0.8% COPD



30-Day Nursing Home CMS Quality Measures

- 21% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 9% of short stay participants were re-hospitalized for any reason
- 3% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*465 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program