



**Rehospitalization Intervention  
Program Objectives:**

- . Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- . Initiate Respiratory Therapy into the Plan of Care
- . Utilize Prevention Strategies

**The U.S. Average for 30-Day  
Re-Hospitalization Rates\*:**

- . 16.7% for Pneumonia
- . 21.7% for CHF
- . 19.6% for COPD

**PEL/VIP Whole Company**  
Cardio-Pulmonary Participant Report  
May 2019


\*medicare.gov/hospitalcompare

**Results Summary: 1252 Total Participants**

Participant Breakdown: Pneumonia - 554 High Risk, 97 Low Risk; CHF - 837 High Risk, 47 Low Risk; COPD - 476 High Risk, 28 Low Risk

Re-Hospitalization Results

1.4% Pneumonia  
0.6% CHF  
1.2% COPD



30-Day Nursing Home CMS Quality Measures

- 22% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 12% of short stay participants were re-hospitalized for any reason
- 3% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*560 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program