



**Rehospitalization Intervention Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies

**The U.S. Average for 30-Day Re-Hospitalization Rates\*:**

- 16.7% for Pneumonia
- 21.7% for CHF
- 19.6% for COPD

**PEL/VIP Whole Company**  
 Cardio-Pulmonary Participant Report  
 August 2019


\*medicare.gov/hospitalcompare

**Results Summary: 1305 Total Participants**

Participant Breakdown: Pneumonia - 555 High Risk, 66 Low Risk; CHF - 837 High Risk, 55 Low Risk; COPD - 551 High Risk, 9 Low Risk

Re-Hospitalization Results

1.0% Pneumonia  
 0.7% CHF  
 0.2% COPD



30-Day Nursing Home CMS Quality Measures

- 24% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 9% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*450 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program