



**Rehospitalization Intervention  
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies

**The U.S. Average for 30-Day  
Re-Hospitalization Rates\*:**

- 16.7% for Pneumonia
- 21.7% for CHF
- 19.6% for COPD

**PEL/VIP Whole Company**  
Cardio-Pulmonary Participant Report  
February 2020


\*medicare.gov/hospitalcompare

**Results Summary: 1661 Total Participants**

Participant Breakdown: Pneumonia - 721 High Risk, 166 Low Risk; CHF - 992 High Risk, 101 Low Risk; COPD - 768 High Risk, 14 Low Risk

Re-Hospitalization Results

0.2% Pneumonia  
0.8% CHF  
0.4% COPD



30-Day Nursing Home CMS Quality Measures

- 21% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 9% of short stay participants were re-hospitalized for any reason
- 3% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*441 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program