



**Rehospitalization Intervention
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- 16.7% for Pneumonia
- 21.7% for CHF
- 19.6% for COPD

PEL/VIP Whole Company
Cardio-Pulmonary Participant Report
March 2020

*medicare.gov/hospitalcompare

Results Summary: 1496 Total Participants


Participant Breakdown: Pneumonia - 608 High Risk, 123 Low Risk; CHF - 885 High Risk, 65 Low Risk; COPD - 651 High Risk, 10 Low Risk

Re-Hospitalization Results

1.0% Pneumonia

0.5% CHF

0.6% COPD



30-Day Nursing Home CMS Quality Measures

- 23% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 9% of short stay participants were re-hospitalized for any reason
- 3% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*415 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program