



**Rehospitalization Intervention  
Program Objectives:**

- Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- Initiate Respiratory Therapy into the Plan of Care
- Utilize Prevention Strategies

**The U.S. Average for 30-Day  
Re-Hospitalization Rates\*:**

- 16.7% for Pneumonia
- 21.7% for CHF
- 19.6% for COPD

**PEL/VIP Whole Company**  
Cardio-Pulmonary Participant Report  
June 2020


\*medicare.gov/hospitalcompare

**Results Summary: 1082 Total Participants**

Participant Breakdown: Pneumonia - 519 High Risk, 79 Low Risk; CHF - 649 High Risk, 46 Low Risk; COPD - 396 High Risk, 2 Low Risk

Re-Hospitalization Results

0.3% Pneumonia  
0.6% CHF  
0.0% COPD



30-Day Nursing Home CMS Quality Measures

- 21% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 5% of short stay participants were re-hospitalized for any reason
- 1% of short stay participants were re-hospitalized for a cardio-pulmonary reason

\*473 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program