



**Rehospitalization Intervention
Program Objectives:**

- . Reduce 30-Day rehospitalizations for Pneumonia, CHF & COPD
- . Initiate Respiratory Therapy into the Plan of Care
- . Utilize Prevention Strategies

**The U.S. Average for 30-Day
Re-Hospitalization Rates*:**

- . 16.7% for Pneumonia
- . 21.7% for CHF
- . 19.6% for COPD

PEL/VIP Whole Company
Cardio-Pulmonary Participant Report
November 2020


*medicare.gov/hospitalcompare

Results Summary: 1198 Total Participants

Participant Breakdown: Pneumonia - 444 High Risk, 122 Low Risk; CHF - 609 High Risk, 64 Low Risk; COPD - 385 High Risk, 30 Low Risk

Re-Hospitalization Results

0.4% Pneumonia
0.1% CHF
0.2% COPD



30-Day Nursing Home CMS Quality Measures

- 21% of short stay participants were successfully discharged to the community
- 0% of short stay participants had an outpatient emergency department visit
- 8% of short stay participants were re-hospitalized for any reason
- 2% of short stay participants were re-hospitalized for a cardio-pulmonary reason

*400 other participants were able to benefit from the use of Respiratory Services outside of the Cardio-Pulmonary Program